



POSTER PRESENTATION

Open Access

Deregulation of IL-1 β axis in peripheral blood mononuclear cells from patients with Chronic Recurrent Multifocal Osteomyelitis

Roberta Scianaro^{1*}, Antonella Insalaco¹, Giusi Prencipe¹, Luisa Bracci-Laudiero^{1,2}, Anna Maria Teti³, Fabrizio De Benedetti¹

From 18th Pediatric Rheumatology European Society (PReS) Congress
Bruges, Belgium. 14-18 September 2011

Background

Chronic recurrent multifocal osteomyelitis (CRMO) is a rare childhood disease presenting with bone pain, fever and multifocal osteolytic bone lesions. CRMO is considered an autoinflammatory disease but its aetiology and pathogenesis are still unknown. Autoinflammatory diseases are usually characterized by abnormalities in the regulation of inflammasomes, macromolecular complexes involved in the processing and in the release of IL-1 β .

Aim

To investigate the inflammasome response in CRMO patients.

Methods

In this study, 10 CRMO patients (mean age 10.7 ± 5.4 years) and 9 age-matched controls were enrolled. The basal mRNA expression of inflammatory cytokines such as IL-1 β , IL-6 and TNF- α and of inflammasome components such as ASC, NLRP3 and CASP-1 were evaluated by quantitative real-time PCR in peripheral blood mononuclear cells (PBMCs). IL-1 β levels released in conditioned media after LPS and ATP treatment of PBMCs were measured by ELISA.

Results

mRNA levels of IL-1 β and IL-6 were significantly higher in PBMCs from CRMO patients than in healthy donors, with a ten-fold average increase in IL1 β transcript ($p=0.0001$) and a five-fold average increase of IL-6

transcript ($p=0.001$). mRNA levels of inflammasome components were comparable between the two groups. When PBMCs were stimulated in vitro with LPS alone, we observed higher levels of IL-1 β in the conditioned media of PBMCs obtained from CRMO patients than in those obtained from controls ($p=0.01$). Controls cells released high IL1 β levels only after stimulation with LPS and ATP with no difference with PBMCs from CRMO patients.

Conclusion

Our data suggest that an abnormal regulation of IL-1 β axis characterize PBMCs of CRMO patients.

Author details

¹IRCCS Ospedale Pediatrico Bambino Gesù, Rome, Italy. ²Istituto di Farmacologia Traslazionale, CNR, Rome, Italy. ³Dipartimento di Medicina Sperimentale, Università dell'Aquila, Italy.

Published: 14 September 2011

doi:10.1186/1546-0096-9-S1-P307

Cite this article as: Scianaro et al.: Deregulation of IL-1 β axis in peripheral blood mononuclear cells from patients with Chronic Recurrent Multifocal Osteomyelitis. *Pediatric Rheumatology* 2011 **9**(Suppl 1):P307.

¹IRCCS Ospedale Pediatrico Bambino Gesù, Rome, Italy
Full list of author information is available at the end of the article